

MODEL-THEORETICAL AND TOPOLOGICAL PROPERTIES OF FAMILIES OF THEORIES

Nurlan Markhabatov

Novosibirsk State Technical University, Novosibirsk, Russia

14TH INTERNATIONAL SUMMER SCHOOL-CONFERENCE ON THE OCCASION OF
THE 75TH BIRTHDAY OF PROFESSOR BRUNO POIZAT

ERLAGOL, JUNE 23-29, 2021

- **Object of study:** The main objects of study in the dissertation are complete, incomplete and pseudofinite theories.
- **Objective:** Investigation of model-theoretical and topological properties of families of theories.

- **Object of study:** The main objects of study in the dissertation are complete, incomplete and pseudofinite theories.
- **Objective:** Investigation of model-theoretical and topological properties of families of theories.

Families of complete theories. Bibliography

- S. V. Sudoplatov, "Combinations of structures", Известия Иркутского государственного университета. Серия Математика, 24 (2018), 82–101
- S. V. Sudoplatov, "Closures and generating sets related to combinations of structures", Известия Иркутского государственного университета. Серия Математика, 16 (2016), 131–144
- S. V. Sudoplatov, "Families of language uniform theories and their generating sets", Известия Иркутского государственного университета. Серия Математика, 17 (2016), 62–76
- S. V. Sudoplatov, "Combinations related to classes of finite and countably categorical structures and their theories", Сиб. электрон. матем. изв., 14 (2017), 135–150
- S. V. Sudoplatov, "Relative e-spectra and relative closures for families of theories", Сиб. электрон. матем. изв., 14 (2017), 296–307
- S. V. Sudoplatov, "On semilattices and lattices for families of theories", Сиб. электрон. матем. изв., 14 (2017), 980–985
- S. V. Sudoplatov, "Approximations of theories", Сиб. электрон. матем. изв., 17 (2020), 715–725
- S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019
- Ю. Л. Ершов, Топология для дискретной математики / Ю.Л. Ершов; отв. ред. В.Л. Селиванов; Рос. акад. наук, Сиб. отд-ние, Ин-т математики им. С.Л. Соболева. – Новосибирск: Изд-во СО РАН, 2020. – 333 с.

We consider the families \mathcal{T} of complete first-order theories of the language $\Sigma = \Sigma(\mathcal{T})$.

- **Definition.**¹ We denote by \mathcal{T}_φ the set $\{T \in \mathcal{T} \mid \varphi \in T\}$.
- Any set \mathcal{T}_φ is called *φ -neighborhood* or just *neighborhood* for \mathcal{T} or a *(φ -)definable* subset of \mathcal{T} .
- The set \mathcal{T}_φ is also called *definable (by a formula or sentence φ)* with respect to \mathcal{T} , or *s-definable*.

¹Sudoplatov, S. V. Closures and generating sets related to combinations of structures / S. V. Sudoplatov // The Bulletin of Irkutsk State University. Series Mathematics. — 2016. — Vol. 16. — P. 131—144.

We consider the families \mathcal{T} of complete first-order theories of the language $\Sigma = \Sigma(\mathcal{T})$.

- **Definition.**¹ We denote by \mathcal{T}_φ the set $\{T \in \mathcal{T} \mid \varphi \in T\}$.
- Any set \mathcal{T}_φ is called φ -neighborhood or just *neighborhood* for \mathcal{T} or a $(\varphi$ -)definable subset of \mathcal{T} .
- The set \mathcal{T}_φ is also called *definable (by a formula or sentence φ)* with respect to \mathcal{T} , or *s-definable*.

¹Sudoplatov, S. V. Closures and generating sets related to combinations of structures / S. V. Sudoplatov // The Bulletin of Irkutsk State University. Series Mathematics. — 2016. — Vol. 16. — P. 131—144.

We consider the families \mathcal{T} of complete first-order theories of the language $\Sigma = \Sigma(\mathcal{T})$.

- **Definition.**¹ We denote by \mathcal{T}_φ the set $\{T \in \mathcal{T} \mid \varphi \in T\}$.
- Any set \mathcal{T}_φ is called φ -neighborhood or just *neighborhood* for \mathcal{T} or a $(\varphi$ -)definable subset of \mathcal{T} .
- The set \mathcal{T}_φ is also called *definable* (by a formula or sentence φ) with respect to \mathcal{T} , or *s-definable*.

¹Sudoplatov, S. V. Closures and generating sets related to combinations of structures / S. V. Sudoplatov // The Bulletin of Irkutsk State University. Series Mathematics. — 2016. — Vol. 16. — P. 131—144.

- **Definition.**² For the empty family \mathcal{T} we put the rank $\text{RS}(\mathcal{T}) = -1$, for finite nonempty families \mathcal{T} we put $\text{RS}(\mathcal{T}) = 0$, and $\text{RS}(\mathcal{T}) \geq 1$ for infinite families \mathcal{T} .
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\text{RS}(\mathcal{T}) \geq \alpha$ if there are pairwise inconsistent $\Sigma(\mathcal{T})$ -sentences φ_n , $n \in \omega$, such that $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal, then $\text{RS}(\mathcal{T}) \geq \alpha$ if $\text{RS}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\text{RS}(\mathcal{T}) = \alpha$ if $\text{RS}(\mathcal{T}) \geq \alpha$ and $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\text{RS}(\mathcal{T}) \geq \alpha$ for any α , we put $\text{RS}(\mathcal{T}) = \infty$.

²S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**² For the empty family \mathcal{T} we put the rank $\text{RS}(\mathcal{T}) = -1$, for finite nonempty families \mathcal{T} we put $\text{RS}(\mathcal{T}) = 0$, and $\text{RS}(\mathcal{T}) \geq 1$ for infinite families \mathcal{T} .
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\text{RS}(\mathcal{T}) \geq \alpha$ if there are pairwise inconsistent $\Sigma(\mathcal{T})$ -sentences φ_n , $n \in \omega$, such that $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal, then $\text{RS}(\mathcal{T}) \geq \alpha$ if $\text{RS}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\text{RS}(\mathcal{T}) = \alpha$ if $\text{RS}(\mathcal{T}) \geq \alpha$ and $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\text{RS}(\mathcal{T}) \geq \alpha$ for any α , we put $\text{RS}(\mathcal{T}) = \infty$.

²S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**² For the empty family \mathcal{T} we put the rank $\text{RS}(\mathcal{T}) = -1$, for finite nonempty families \mathcal{T} we put $\text{RS}(\mathcal{T}) = 0$, and $\text{RS}(\mathcal{T}) \geq 1$ for infinite families \mathcal{T} .
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\text{RS}(\mathcal{T}) \geq \alpha$ if there are pairwise inconsistent $\Sigma(\mathcal{T})$ -sentences φ_n , $n \in \omega$, such that $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal, then $\text{RS}(\mathcal{T}) \geq \alpha$ if $\text{RS}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\text{RS}(\mathcal{T}) = \alpha$ if $\text{RS}(\mathcal{T}) \geq \alpha$ and $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\text{RS}(\mathcal{T}) \geq \alpha$ for any α , we put $\text{RS}(\mathcal{T}) = \infty$.

²S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**² For the empty family \mathcal{T} we put the rank $\text{RS}(\mathcal{T}) = -1$, for finite nonempty families \mathcal{T} we put $\text{RS}(\mathcal{T}) = 0$, and $\text{RS}(\mathcal{T}) \geq 1$ for infinite families \mathcal{T} .
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\text{RS}(\mathcal{T}) \geq \alpha$ if there are pairwise inconsistent $\Sigma(\mathcal{T})$ -sentences φ_n , $n \in \omega$, such that $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal, then $\text{RS}(\mathcal{T}) \geq \alpha$ if $\text{RS}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\text{RS}(\mathcal{T}) = \alpha$ if $\text{RS}(\mathcal{T}) \geq \alpha$ and $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\text{RS}(\mathcal{T}) \geq \alpha$ for any α , we put $\text{RS}(\mathcal{T}) = \infty$.

²S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**² For the empty family \mathcal{T} we put the rank $\text{RS}(\mathcal{T}) = -1$, for finite nonempty families \mathcal{T} we put $\text{RS}(\mathcal{T}) = 0$, and $\text{RS}(\mathcal{T}) \geq 1$ for infinite families \mathcal{T} .
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\text{RS}(\mathcal{T}) \geq \alpha$ if there are pairwise inconsistent $\Sigma(\mathcal{T})$ -sentences φ_n , $n \in \omega$, such that $\text{RS}(\mathcal{T}_{\varphi_n}) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal, then $\text{RS}(\mathcal{T}) \geq \alpha$ if $\text{RS}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\text{RS}(\mathcal{T}) = \alpha$ if $\text{RS}(\mathcal{T}) \geq \alpha$ and $\text{RS}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\text{RS}(\mathcal{T}) \geq \alpha$ for any α , we put $\text{RS}(\mathcal{T}) = \infty$.

²S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**³ A family \mathcal{T} is called *e-totally transcendental*, or *totally transcendental*, if $\text{RS}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is totally transcendental, with $\text{RS}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\text{ds}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise inconsistent sentences φ_i such that $\text{RS}(\mathcal{T}_{\varphi_i}) = \alpha$.

³S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**³ A family \mathcal{T} is called *e-totally transcendental*, or *totally transcendental*, if $\text{RS}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is totally transcendental, with $\text{RS}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\text{ds}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise inconsistent sentences φ_i such that $\text{RS}(\mathcal{T}_{\varphi_i}) = \alpha$.

³S. V. Sudoplatov, Ranks for families of theories and their spectra, Lobachevskii Journal of Mathematics (to appear). arXiv:1901.08464v1[math.LO]. 2019

- **Definition.**⁴ Let \mathcal{T} be a family of theories and T be a theory, $T \notin \mathcal{T}$. The theory T is called \mathcal{T} -approximated, or approximated by \mathcal{T} , or \mathcal{T} -approximable, or a pseudo- \mathcal{T} -theory, if for any formula $\varphi \in T$ there is $T' \in \mathcal{T}$ such that $\varphi \in T'$.
- If T is \mathcal{T} -approximated then \mathcal{T} is called an approximating family for T , theories $T' \in \mathcal{T}$ are approximations for T , and T is an accumulation point for \mathcal{T} .
- An approximating family \mathcal{T} is called e -minimal if for any sentence $\varphi \in \Sigma(\mathcal{T})$, \mathcal{T}_φ is finite or $\mathcal{T}_{\neg\varphi}$ is finite.

⁴S. V. Sudoplatov, “Approximations of theories”, Сиб. электрон. матем. изв., 17 (2020), 715–725

- **Definition.**⁴ Let \mathcal{T} be a family of theories and T be a theory, $T \notin \mathcal{T}$. The theory T is called \mathcal{T} -approximated, or approximated by \mathcal{T} , or \mathcal{T} -approximable, or a pseudo- \mathcal{T} -theory, if for any formula $\varphi \in T$ there is $T' \in \mathcal{T}$ such that $\varphi \in T'$.
- If T is \mathcal{T} -approximated then \mathcal{T} is called an approximating family for T , theories $T' \in \mathcal{T}$ are approximations for T , and T is an accumulation point for \mathcal{T} .
- An approximating family \mathcal{T} is called *e-minimal* if for any sentence $\varphi \in \Sigma(\mathcal{T})$, \mathcal{T}_φ is finite or $\mathcal{T}_{\neg\varphi}$ is finite.

⁴S. V. Sudoplatov, “Approximations of theories”, Сиб. электрон. матем. изв., 17 (2020), 715–725

- **Definition.**⁴ Let \mathcal{T} be a family of theories and T be a theory, $T \notin \mathcal{T}$. The theory T is called \mathcal{T} -approximated, or approximated by \mathcal{T} , or \mathcal{T} -approximable, or a pseudo- \mathcal{T} -theory, if for any formula $\varphi \in T$ there is $T' \in \mathcal{T}$ such that $\varphi \in T'$.
- If T is \mathcal{T} -approximated then \mathcal{T} is called an approximating family for T , theories $T' \in \mathcal{T}$ are approximations for T , and T is an accumulation point for \mathcal{T} .
- An approximating family \mathcal{T} is called e -minimal if for any sentence $\varphi \in \Sigma(\mathcal{T})$, \mathcal{T}_φ is finite or $\mathcal{T}_{\neg\varphi}$ is finite.

⁴S. V. Sudoplatov, “Approximations of theories”, Сиб. электрон. матем. изв., 17 (2020), 715–725

- Recall that the E -closure $Cl_E(\mathcal{T})$ ⁵ for the family \mathcal{T} of complete theories is characterized by the following proposition.
- **Proposition.** *Let \mathcal{T} be a family of complete theories of the language Σ . Then $Cl_E(\mathcal{T}) = \mathcal{T}$ for finite \mathcal{T} and for infinite \mathcal{T} , the theory T belongs to $Cl_E(\mathcal{T})$ if and only if T is a complete theory of the language Σ and $T \in \mathcal{T}$, or $T \notin \mathcal{T}$ and for of any formula $\varphi \in T$ the set \mathcal{T}_φ is infinite.*

⁵Sudoplatov, S. V. Closures and generating sets related to combinations of structures / S. V. Sudoplatov // The Bulletin of Irkutsk State University. Series Mathematics. — 2016. — Vol. 16. — P. 131—144

- Recall that the E -closure $Cl_E(\mathcal{T})$ ⁵ for the family \mathcal{T} of complete theories is characterized by the following proposition.
- Proposition.** *Let \mathcal{T} be a family of complete theories of the language Σ . Then $Cl_E(\mathcal{T}) = \mathcal{T}$ for finite \mathcal{T} and for infinite \mathcal{T} , the theory T belongs to $Cl_E(\mathcal{T})$ if and only if T is a complete theory of the language Σ and $T \in \mathcal{T}$, or $T \notin \mathcal{T}$ and for of any formula $\varphi \in T$ the set \mathcal{T}_φ is infinite.*

⁵Sudoplatov, S. V. Closures and generating sets related to combinations of structures / S. V. Sudoplatov // The Bulletin of Irkutsk State University. Series Mathematics. — 2016. — Vol. 16. — P. 131—144

Regularly closed, pseudo-algebraically closed, quasifinite fields

It should be noted are actively developing in recent area of study **pseudofinite** structures.

- **F is regularly closed**,⁶ *if every absolutely irreducible variety V over F has a point defined over F .*
- Ax fields, Σ -fields, Pseudo-algebraically closed fields⁷
- **F is quasifinite**, if it is perfect and has exactly one extension of each degree (in the fixed algebraic closure \tilde{F}).

⁶Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁷ Ax , J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

Regularly closed, pseudo-algebraically closed, quasifinite fields

It should be noted are actively developing in recent area of study **pseudofinite** structures.

- **F is regularly closed**,⁶ *if every absolutely irreducible variety V over F has a point defined over F .*
- **Ax fields, Σ -fields, Pseudo-algebraically closed fields**⁷
- **F is quasifinite**, if it is perfect and has exactly one extension of each degree (in the fixed algebraic closure \tilde{F}).

⁶Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁷ Ax , J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

Regularly closed, pseudo-algebraically closed, quasifinite fields

It should be noted are actively developing in recent area of study **pseudofinite** structures.

- **F is regularly closed**,⁶ if every absolutely irreducible variety V over F has a point defined over F .
- A_x fields, Σ -fields, **Pseudo-algebraically closed fields**⁷
- **F is quasifinite**, if it is perfect and has exactly one extension of each degree (in the fixed algebraic closure \tilde{F}).

⁶Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁷ A_x , J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

- Ершов, Ю. Л. Регулярно замкнутые поля. Докл. АН СССР. 1980. Т. 251, No 4. С. 783—785.
- Ершов, Ю. Л. Неразрешимость регулярно замкнутых полей. Алгебра и логика. 1981. Т. 20, No 4. С. 389—394.
- Ершов, Ю. Л. Алгебраические свойства регулярно замкнутых полей. Тр. МИАН СССР. 1981. Т. 158. С. 80—86.
- G. Cherlin, L. van den Dries, A. Macintyre. The elementary theory of regularly closed fields. preprint.

- M. Fried, G. Sacerdote, Solving diophantihe problems over all residue class fields. Ann. Math. 1976. Vol. 104. P. 203—233.
- Jarden, M. The elementary theory of ω -free Ax fields. Invent.Math. 1976. Vol. 38. P. 187—206.
- M. Jarden, S. Shelah, Pseudo-algebraically closed fields over rational function fields. Proceedings of the American Mathematical Society. 1983. Vol. 87, no. 2. P. 223—223.
- Frey, G. Pseudo algebraically closed fields with non-archimedian real valuations. Journal of Algebra. 1973. Vol. 26. P. 202—207.

- Wheeler W. H., Jarden M., Model-complete theories of e -free Ax fields, The Journal of Symbolic Logic. 1983. Vol. 48, no. 4. P. 1125—1129.
- Wheeler, W. H. Model-complete theories of pseudo-algebraically closed fields. Ann.Math. Log. 1979. Vol. 17, no. 3. P. 205—226.
- Z. Chatzidakis, E. Hrushovski, Perfect pseudo-algebraically closed fields are algebraically bounded. Journal of Algebra. 2004. Vol. 271, no. 2. P. 627—637.
- A. Pillay, D. Polkowska, On PAC and bounded substructures of a stable structure. The Journal of Symbolic Logic. 2006. Vol. 71, no. 2. P. 460—472.

Pseudofinite theories

- **Theorem.** (Ax, 1968) The field F is *pseudofinite* if and only if all of the following conditions are met:
 - (i) F is perfect;
 - (ii) F is quasifinite;
 - (iii) F is pseudo-algebraically closed (PAC)^{8 9}.
- **Proposition.** (Ax, 1968) We fix the language Σ and the Σ -structure \mathcal{M} . Then the following are equivalent:
 - 1. The Σ -structure \mathcal{M} is pseudofinite;
 - 2. $\mathcal{M} \models T_f$, where T_f is the general theory of all finite Σ -structures;
 - 3. \mathcal{M} is elementarily equivalent to an ultraproduct of finite Σ -structures.
- **Definition.**¹⁰ A Σ -structure \mathcal{M} is called *pseudofinite* if, for all Σ -sentences φ , it follows from $\mathcal{M} \models \varphi$ that there is a finite \mathcal{M}_0 such that $\mathcal{M}_0 \models \varphi$.

⁸Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁹Ax, J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

¹⁰Ax, J. The elementary theory of finite fields, Ann. Math. 1968, Vol. 88, P. 239—271

Pseudofinite theories

- **Theorem.** (Ax, 1968) The field F is *pseudofinite* if and only if all of the following conditions are met:
 - (i) F is perfect;
 - (ii) F is quasifinite;
 - (iii) F is pseudo-algebraically closed (PAC)^{8 9}.
- **Proposition.** (Ax, 1968) We fix the language Σ and the Σ -structure \mathcal{M} . Then the following are equivalent:
 - 1. The Σ -structure \mathcal{M} is pseudofinite;
 - 2. $\mathcal{M} \models T_f$, where T_f is the general theory of all finite Σ -structures;
 - 3. \mathcal{M} is elementarily equivalent to an ultraproduct of finite Σ -structures.
- **Definition.**¹⁰ A Σ -structure \mathcal{M} is called *pseudofinite* if, for all Σ -sentences φ , it follows from $\mathcal{M} \models \varphi$ that there is a finite \mathcal{M}_0 such that $\mathcal{M}_0 \models \varphi$.

⁸Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁹Ax, J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

¹⁰Ax, J. The elementary theory of finite fields, Ann. Math. 1968, Vol. 88, P. 239—271

- **Theorem.** (Ax, 1968) The field F is *pseudofinite* if and only if all of the following conditions are met:
 - (i) F is perfect;
 - (ii) F is quasifinite;
 - (iii) F is pseudo-algebraically closed (PAC)^{8 9}.
- **Proposition.** (Ax, 1968) We fix the language Σ and the Σ -structure \mathcal{M} . Then the following are equivalent:
 - 1. The Σ -structure \mathcal{M} is pseudofinite;
 - 2. $\mathcal{M} \models T_f$, where T_f is the general theory of all finite Σ -structures;
 - 3. \mathcal{M} is elementarily equivalent to an ultraproduct of finite Σ -structures.
- **Definition.**¹⁰ A Σ -structure \mathcal{M} is called **pseudofinite** if, for all Σ -sentences φ , it follows from $\mathcal{M} \models \varphi$ that there is a finite \mathcal{M}_0 such that $\mathcal{M}_0 \models \varphi$.

⁸Ершов, Ю. Л. О полях с разрешимой теорией. Докл. АН СССР. 1967. Т. 174, No 1. С. 19—20.

⁹Ax, J. Solving diophantine problems modulo every prime. Ann. Math. 1967. Vol. 85, no. 2. P. 161—183.

¹⁰Ax, J. The elementary theory of finite fields, Ann. Math. 1968. Vol. 88. P. 239—271.

Some examples

- DLO theory is not pseudofinite
- The Abelian group $(\mathbb{Z}, +)$ is not pseudofinite.
- The theory of algebraically closed fields is not pseudofinite

Some examples

- DLO theory is not pseudofinite
- The Abelian group $(\mathbb{Z}, +)$ is not pseudofinite.
- The theory of algebraically closed fields is not pseudofinite

Some examples

- DLO theory is not pseudofinite
- The Abelian group $(\mathbb{Z}, +)$ is not pseudofinite.
- The theory of algebraically closed fields is not pseudofinite

Pseudofinite structures (fields, groups, rings). Bibliography

Pseudofinite fields:

- Cherlin, G. Finite Structures with Few Types / G. Cherlin, E. Hrushovski // Annals of Math. Studies, Princeton University Press. 2003. Vol. 152.
- Chatzidakis, Z. Model theory of finite fields and pseudofinite fields / Z. Chatzidakis // Ann. Pure Appl. Logic. 1997. Vol. 88. P. 95—108.

Pseudofinite groups:

- Wilson, J. On pseudofinite simple groups. J. Lond. Math. Soc. 1995. Vol. 51, no. 2. P. 471—490.
- Point, F. Ultraproducts and Chevalley groups. Arch. Math. Logic. 1999. Vol. 38. P. 355—372.
- Macpherson, H. Stable pseudofinite groups / H. Macpherson, K. Tent // J. Algebra. 2007. Vol. 312. P. 550—561.
- Macpherson, D. Model theory of finite and pseudofinite groups. Arch. Math. Logic. 2018. Vol. 57. P. 159—184.
- Pavlyuk, I. I. Approximations for Theories of Abelian Groups / I. I. Pavlyuk, S. V. Sudoplatov // Mathematics and Statistics. 2020. Vol. 8, no. 2. P. 220—224.

Pseudofinite rings:

- Bello-Aguirre, R. Model Theory of Finite and Pseudofinite Rings : PhD thesis / Bello-Aguirre R. University of Leeds, 2016.

Basic results on pseudofinite theories

- **Theorem.** ¹¹ The theory of any pseudo-algebraically closed (PAC) field that is not separably closed has the independence property. (That is, the pseudo-finite theory is unstable)
- **Theorem.** ¹² Totally categorical theories (and more generally ω -categorical ω -stable theories¹³) are pseudofinite.

¹¹Duret, J. Les corps pseudo-finis ont la propriete d'indpendance. C. R. Acad. Sei. Paris Ser. 1980. Vol. 290. P. 981—903.

¹²Zilber, B. On the problem of finite axiomatizability for theories categorical in all infinite powers (Russian) in: B. Bajjanov, ed., Investigations in Theoretical Programming (Alma Ata 1981). 1981. P. 69—75.

¹³Cherlin G., Harrington L. and Lachlan A. H., \aleph_0 -categorical, \aleph_0 -stable structures, Ann. Pure Appl. Logic 28 (1985) 103–135.

Basic results on pseudofinite theories

- **Theorem.** ¹¹ The theory of any pseudo-algebraically closed (PAC) field that is not separably closed has the independence property. (That is, the pseudo-finite theory is unstable)
- **Theorem.** ¹² Totally categorical theories (and more generally ω -categorical ω -stable theories¹³) are pseudofinite.

¹¹Duret, J. Les corps pseudo-finis ont la propriete d'indpendance. C. R. Acad. Sei. Paris Ser. 1980. Vol. 290. P. 981—903.

¹²Zilber, B. On the problem of finite axiomatizability for theories categorical in all infinite powers (Russian) in: B. Baijanov, ed., Investigations in Theoretical Programming (Alma Ata 1981). 1981. P. 69—75.

¹³Cherlin G., Harrington L. and Lachlan A. H., \aleph_0 -categorical, \aleph_0 -stable structures, Ann. Pure Appl. Logic 28 (1985) 103–135.

Ranks for families of theories of arbitrary languages(with S.V.Sudoplatov)¹⁴

Let Σ be a language. If Σ is relational, then by \mathcal{T}_Σ we denote the family of all theories of the language Σ .

- **Theorem 1.** For any language Σ the family \mathcal{T}_Σ is e-minimal if and only if $\Sigma = \emptyset$ or Σ consists of one constant symbol.
- **Theorem 2.** For any language Σ either $\text{RS}(\mathcal{T}_\Sigma)$ is finite, if Σ consists of finitely many 0-ary and unary predicates, and finitely many constant symbols, or $\text{RS}(\mathcal{T}_\Sigma) = \infty$, otherwise.

¹⁴Markhabatov, N. D. Ranks for families of all theories of given languages /N. D. Markhabatov, S. V. Sudoplatov // Eurasian Mathematical Journal, 2021 (to appear) arXiv:1901.09903v1 [math.LO]. — 2019.

Ranks for families of theories of arbitrary languages(with S.V.Sudoplatov)¹⁴

Let Σ be a language. If Σ is relational, then by \mathcal{T}_Σ we denote the family of all theories of the language Σ .

- **Theorem 1.** For any language Σ the family \mathcal{T}_Σ is e-minimal if and only if $\Sigma = \emptyset$ or Σ consists of one constant symbol.
- **Theorem 2.** For any language Σ either $\text{RS}(\mathcal{T}_\Sigma)$ is finite, if Σ consists of finitely many 0-ary and unary predicates, and finitely many constant symbols, or $\text{RS}(\mathcal{T}_\Sigma) = \infty$, otherwise.

¹⁴Markhabatov, N. D. Ranks for families of all theories of given languages /N. D. Markhabatov, S. V. Sudoplatov // Eurasian Mathematical Journal, 2021 (to appear) arXiv:1901.09903v1 [math.LO]. — 2019.

Calculi for definable families of complete theories (with S.V.Sudoplatov)¹⁵

- **Definition.** For a family \mathcal{T} and sentences φ and ψ we say that φ \mathcal{T} -forces ψ , written $\varphi \vdash_{\mathcal{T}} \psi$ if $\mathcal{T}_{\varphi} \subseteq \mathcal{T}_{\psi}$.
- Definition. If \mathcal{T} is a family of theories and Φ is a set of sentences, then we put $\mathcal{T}_{\Phi} = \bigcap_{\varphi \in \Phi} \mathcal{T}_{\varphi}$ and the set \mathcal{T}_{Φ} is called (type- or diagram-)definable (by the set Φ) with respect to \mathcal{T} , or (diagram-) \mathcal{T} -definable, or simply d -definable.
- If Φ is a singleton $\{\varphi\}$, then $\mathcal{T}_{\varphi} = \mathcal{T}_{\Phi}$ is called s -definable.
- **Proposition 3.** For any sets Φ and Ψ of sentences and a family \mathcal{T} of theories the following conditions are equivalent:
 - (1) $\Phi \vdash_{\mathcal{T}} \Psi$;
 - (2) $\Phi \vdash_{\mathcal{T}_0} \Psi$ for any finite $\mathcal{T}_0 \subseteq \mathcal{T}$;
 - (3) $\Phi \vdash_{\{T\}} \Psi$ for any singleton $\{T\} \subseteq \mathcal{T}$;
 - (4) $\Phi \vdash_{Cl_E(\mathcal{T})} \Psi$.

¹⁵Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Calculi for definable families of complete theories (with S.V.Sudoplatov)¹⁵

- **Definition.** For a family \mathcal{T} and sentences φ and ψ we say that φ \mathcal{T} -forces ψ , written $\varphi \vdash_{\mathcal{T}} \psi$ if $\mathcal{T}_{\varphi} \subseteq \mathcal{T}_{\psi}$.
- **Definition.** If \mathcal{T} is a family of theories and Φ is a set of sentences, then we put $\mathcal{T}_{\Phi} = \bigcap_{\varphi \in \Phi} \mathcal{T}_{\varphi}$ and the set \mathcal{T}_{Φ} is called (*type- or diagram-*)*definable* (by the set Φ) with respect to \mathcal{T} , or (*diagram-*) \mathcal{T} -*definable*, or simply *d-definable*.
- If Φ is a singleton $\{\varphi\}$, then $\mathcal{T}_{\varphi} = \mathcal{T}_{\Phi}$ is called *s-definable*.
- **Proposition 3.** For any sets Φ and Ψ of sentences and a family \mathcal{T} of theories the following conditions are equivalent:
 - (1) $\Phi \vdash_{\mathcal{T}} \Psi$;
 - (2) $\Phi \vdash_{\mathcal{T}_0} \Psi$ for any finite $\mathcal{T}_0 \subseteq \mathcal{T}$;
 - (3) $\Phi \vdash_{\{T\}} \Psi$ for any singleton $\{T\} \subseteq \mathcal{T}$;
 - (4) $\Phi \vdash_{Cl_E(\mathcal{T})} \Psi$.

¹⁵Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Calculi for definable families of complete theories (with S.V.Sudoplatov)¹⁵

- **Definition.** For a family \mathcal{T} and sentences φ and ψ we say that φ \mathcal{T} -forces ψ , written $\varphi \vdash_{\mathcal{T}} \psi$ if $\mathcal{T}_{\varphi} \subseteq \mathcal{T}_{\psi}$.
- **Definition.** If \mathcal{T} is a family of theories and Φ is a set of sentences, then we put $\mathcal{T}_{\Phi} = \bigcap_{\varphi \in \Phi} \mathcal{T}_{\varphi}$ and the set \mathcal{T}_{Φ} is called (*type- or diagram-*)*definable* (by the set Φ) with respect to \mathcal{T} , or (*diagram-*) \mathcal{T} -*definable*, or simply *d-definable*.
- If Φ is a singleton $\{\varphi\}$, then $\mathcal{T}_{\varphi} = \mathcal{T}_{\Phi}$ is called *s-definable*.
- **Proposition 3.** For any sets Φ and Ψ of sentences and a family \mathcal{T} of theories the following conditions are equivalent:
 - (1) $\Phi \vdash_{\mathcal{T}} \Psi$;
 - (2) $\Phi \vdash_{\mathcal{T}_0} \Psi$ for any finite $\mathcal{T}_0 \subseteq \mathcal{T}$;
 - (3) $\Phi \vdash_{\{T\}} \Psi$ for any singleton $\{T\} \subseteq \mathcal{T}$;
 - (4) $\Phi \vdash_{Cl_E(\mathcal{T})} \Psi$.

¹⁵Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Calculi for definable families of complete theories (with S.V.Sudoplatov)¹⁵

- **Definition.** For a family \mathcal{T} and sentences φ and ψ we say that φ \mathcal{T} -forces ψ , written $\varphi \vdash_{\mathcal{T}} \psi$ if $\mathcal{T}_{\varphi} \subseteq \mathcal{T}_{\psi}$.
- **Definition.** If \mathcal{T} is a family of theories and Φ is a set of sentences, then we put $\mathcal{T}_{\Phi} = \bigcap_{\varphi \in \Phi} \mathcal{T}_{\varphi}$ and the set \mathcal{T}_{Φ} is called (type- or diagram-)definable (by the set Φ) with respect to \mathcal{T} , or (diagram-) \mathcal{T} -definable, or simply d -definable.
- If Φ is a singleton $\{\varphi\}$, then $\mathcal{T}_{\varphi} = \mathcal{T}_{\Phi}$ is called s -definable.
- **Proposition 3.** For any sets Φ and Ψ of sentences and a family \mathcal{T} of theories the following conditions are equivalent:
 - (1) $\Phi \vdash_{\mathcal{T}} \Psi$;
 - (2) $\Phi \vdash_{\mathcal{T}_0} \Psi$ for any finite $\mathcal{T}_0 \subseteq \mathcal{T}$;
 - (3) $\Phi \vdash_{\{T\}} \Psi$ for any singleton $\{T\} \subseteq \mathcal{T}$;
 - (4) $\Phi \vdash_{Cl_E(\mathcal{T})} \Psi$.

¹⁵Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Compactness of definable families of complete theories(with S.V.Sudoplatov)¹⁶

- **Definition.** A d -definable set \mathcal{T}_Φ is called \mathcal{T} -consistent if $\mathcal{T}_\Phi \cap \mathcal{T} \neq \emptyset$, and \mathcal{T}_Φ is called *locally \mathcal{T} -consistent* if for any finite $\Phi_0 \subseteq \Phi$, \mathcal{T}_{Φ_0} is \mathcal{T} -consistent.
- **Theorem 4.** (Compactness theorem) *For any nonempty E -closed family \mathcal{T} , every locally \mathcal{T} -consistent d -definable set \mathcal{T}_Φ is \mathcal{T} -consistent.*
- The following theorem gives a criterion of existence of d -definable family which is not s -definable.
- **Theorem 5.** *For any E -closed family \mathcal{T} , there is a d -definable family \mathcal{T}_Φ which is not s -definable if and only if \mathcal{T} is infinite.*

¹⁶Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Compactness of definable families of complete theories(with S.V.Sudoplatov)¹⁶

- **Definition.** A d -definable set \mathcal{T}_Φ is called \mathcal{T} -consistent if $\mathcal{T}_\Phi \cap \mathcal{T} \neq \emptyset$, and \mathcal{T}_Φ is called *locally \mathcal{T} -consistent* if for any finite $\Phi_0 \subseteq \Phi$, \mathcal{T}_{Φ_0} is \mathcal{T} -consistent.
- **Theorem 4.** (Compactness theorem) *For any nonempty E -closed family \mathcal{T} , every locally \mathcal{T} -consistent d -definable set \mathcal{T}_Φ is \mathcal{T} -consistent.*
- The following theorem gives a criterion of existence of d -definable family which is not s -definable.
- **Theorem 5.** *For any E -closed family \mathcal{T} , there is a d -definable family \mathcal{T}_Φ which is not s -definable if and only if \mathcal{T} is infinite.*

¹⁶Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Compactness of definable families of complete theories(with S.V.Sudoplatov)¹⁶

- **Definition.** A d -definable set \mathcal{T}_Φ is called \mathcal{T} -consistent if $\mathcal{T}_\Phi \cap \mathcal{T} \neq \emptyset$, and \mathcal{T}_Φ is called *locally \mathcal{T} -consistent* if for any finite $\Phi_0 \subseteq \Phi$, \mathcal{T}_{Φ_0} is \mathcal{T} -consistent.
- **Theorem 4.** (Compactness theorem) *For any nonempty E -closed family \mathcal{T} , every locally \mathcal{T} -consistent d -definable set \mathcal{T}_Φ is \mathcal{T} -consistent.*
- The following theorem gives a criterion of existence of d -definable family which is not s -definable.
- **Theorem 5.** *For any E -closed family \mathcal{T} , there is a d -definable family \mathcal{T}_Φ which is not s -definable if and only if \mathcal{T} is infinite.*

¹⁶Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Compactness of definable families of complete theories(with S.V.Sudoplatov)¹⁶

- **Definition.** A d -definable set \mathcal{T}_Φ is called \mathcal{T} -consistent if $\mathcal{T}_\Phi \cap \mathcal{T} \neq \emptyset$, and \mathcal{T}_Φ is called *locally \mathcal{T} -consistent* if for any finite $\Phi_0 \subseteq \Phi$, \mathcal{T}_{Φ_0} is \mathcal{T} -consistent.
- **Theorem 4.** (Compactness theorem) *For any nonempty E -closed family \mathcal{T} , every locally \mathcal{T} -consistent d -definable set \mathcal{T}_Φ is \mathcal{T} -consistent.*
- The following theorem gives a criterion of existence of d -definable family which is not s -definable.
- **Theorem 5.** *For any E -closed family \mathcal{T} , there is a d -definable family \mathcal{T}_Φ which is not s -definable if and only if \mathcal{T} is infinite.*

¹⁶Markhabatov, N. D. Definable families of theories, related calculi and ranks / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2020. — T. 17. — C. 700—714.

Algebras for definable subfamilies of complete theories (with S.V.Sudoplatov)¹⁷

- **Theorem 6.** For any nonempty E -closed family \mathcal{T} and $n \in \omega \setminus \{0\}$ the following conditions are equivalent:
 - (1) $\text{RS}(\mathcal{T}) = 1$ and $\text{ds}(\mathcal{T}) = n$;
 - (2) *Boolean algebra $\mathcal{B}_s(\mathcal{T})$ is isomorphic to the direct product n of Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$, each of which is generated by elements of rank $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
 - (3) *the algebra $\mathcal{B}_d(\mathcal{T})$ consists of an infinite number of atomic elements of each rank $\beta < \alpha, \beta \geq 0$ and exactly n atomic elements of rank α , these n atomic elements correspond to nonprincipal ultrafilters with respect to elements of rank $< \alpha$.*

¹⁷Markhabatov, N. D. Algebras for definable families of theories / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2019. — T. 16. — C. 600—608.

Algebras for definable subfamilies of complete theories (with S.V.Sudoplatov)¹⁷

- **Theorem 6.** For any nonempty E -closed family \mathcal{T} and $n \in \omega \setminus \{0\}$ the following conditions are equivalent:
 - (1) $\text{RS}(\mathcal{T}) = 1$ and $\text{ds}(\mathcal{T}) = n$;
 - (2) *Boolean algebra $\mathcal{B}_s(\mathcal{T})$ is isomorphic to the direct product n of Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$, each of which is generated by elements of rank $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
 - (3) *the algebra $\mathcal{B}_d(\mathcal{T})$ consists of an infinite number of atomic elements of each rank $\beta < \alpha, \beta \geq 0$ and exactly n atomic elements of rank α , these n atomic elements correspond to nonprinciple ultrafilters with respect to elements of rank $< \alpha$.*

¹⁷Markhabatov, N. D. Algebras for definable families of theories / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2019. — T. 16. — C. 600—608.

Algebras for definable subfamilies of complete theories(with S.V.Sudoplatov)¹⁷

- **Theorem 6.** For any nonempty E -closed family \mathcal{T} and $n \in \omega \setminus \{0\}$ the following conditions are equivalent:
(1) $\text{RS}(\mathcal{T}) = 1$ and $\text{ds}(\mathcal{T}) = n$;
- (2) *Boolean algebra $\mathcal{B}_s(\mathcal{T})$ is isomorphic to the direct product n of Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$, each of which is generated by elements of rank $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
- (3) *the algebra $\mathcal{B}_d(\mathcal{T})$ consists of an infinite number of atomic elements of each rank $\beta < \alpha, \beta \geq 0$ and exactly n atomic elements of rank α , these n atomic elements correspond to nonprinciple ultrafilters with respect to elements of rank $< \alpha$.*

¹⁷Markhabatov, N. D. Algebras for definable families of theories / N. D. Markhabatov, S. V. Sudoplatov // Siberian Electronic Mathematical Reports. — 2019. — T. 16. — C. 600—608.

Topologies for families of complete theories(with S.V.Sudoplatov)¹⁸

- We consider families \mathcal{T} of consistent first-order theories in languages $\Sigma \subseteq \Sigma(\mathcal{T})$, not necessary complete. By $F(\Sigma)$ we denote the set of all formulas in the language Σ and by $\text{Sent}(\Sigma)$ the set of all sentences in $F(\Sigma)$.
- **Proposition 7.** *Any family \mathcal{T} forms a T_0 -space.*
- **Proposition 8.** *For any family \mathcal{T} the following conditions are equivalent:*
 - (1) \mathcal{T} is a T_1 -space;
 - (2) \mathcal{T} does not contain theories T, T' with $T \subsetneq T'$.
- **Proposition 9.** *A T_1 -space \mathcal{T} is Hausdorff if and only if for any distinct theories $T, T' \in \mathcal{T}$ there are sentences $\varphi \in T, \psi \in T'$ such that $\varphi \wedge \psi$ is \mathcal{T} -inconsistent.*
- **Theorem 10.** *For any ordinals $\alpha \leq \beta$ and natural $m, n \in \omega \setminus \{0\}$, where $m \leq n$ if $\beta = \alpha$, there is a family \mathcal{T} and its Σ' -restriction \mathcal{T}' such that $\text{RS}(\mathcal{T}) = \alpha$, $\text{ds}(\mathcal{T}) = m$, $\text{RS}(\mathcal{T}') = \beta$, $\text{ds}(\mathcal{T}') = n$.*

¹⁸Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Topologies for families of complete theories(with S.V.Sudoplatov)¹⁸

- We consider families \mathcal{T} of consistent first-order theories in languages $\Sigma \subseteq \Sigma(\mathcal{T})$, not necessary complete. By $F(\Sigma)$ we denote the set of all formulas in the language Σ and by $\text{Sent}(\Sigma)$ the set of all sentences in $F(\Sigma)$.
- **Proposition 7.** *Any family \mathcal{T} forms a T_0 -space.*
- **Proposition 8.** *For any family \mathcal{T} the following conditions are equivalent:*
 - (1) \mathcal{T} is a T_1 -space;
 - (2) \mathcal{T} does not contain theories T, T' with $T \subsetneq T'$.
- **Proposition 9.** *A T_1 -space \mathcal{T} is Hausdorff if and only if for any distinct theories $T, T' \in \mathcal{T}$ there are sentences $\varphi \in T, \psi \in T'$ such that $\varphi \wedge \psi$ is \mathcal{T} -inconsistent.*
- **Theorem 10.** *For any ordinals $\alpha \leq \beta$ and natural $m, n \in \omega \setminus \{0\}$, where $m \leq n$ if $\beta = \alpha$, there is a family \mathcal{T} and its Σ' -restriction \mathcal{T}' such that $\text{RS}(\mathcal{T}) = \alpha$, $\text{ds}(\mathcal{T}) = m$, $\text{RS}(\mathcal{T}') = \beta$, $\text{ds}(\mathcal{T}') = n$.*

¹⁸Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Topologies for families of complete theories(with S.V.Sudoplatov)¹⁸

- We consider families \mathcal{T} of consistent first-order theories in languages $\Sigma \subseteq \Sigma(\mathcal{T})$, not necessary complete. By $F(\Sigma)$ we denote the set of all formulas in the language Σ and by $\text{Sent}(\Sigma)$ the set of all sentences in $F(\Sigma)$.
- **Proposition 7.** Any family \mathcal{T} forms a T_0 -space.
- **Proposition 8.** For any family \mathcal{T} the following conditions are equivalent:
 - (1) \mathcal{T} is a T_1 -space;
 - (2) \mathcal{T} does not contain theories T, T' with $T \subsetneq T'$.
- **Proposition 9.** A T_1 -space \mathcal{T} is Hausdorff if and only if for any distinct theories $T, T' \in \mathcal{T}$ there are sentences $\varphi \in T, \psi \in T'$ such that $\varphi \wedge \psi$ is \mathcal{T} -inconsistent.
- **Theorem 10.** For any ordinals $\alpha \leq \beta$ and natural $m, n \in \omega \setminus \{0\}$, where $m \leq n$ if $\beta = \alpha$, there is a family \mathcal{T} and its Σ' -restriction \mathcal{T}' such that $\text{RS}(\mathcal{T}) = \alpha$, $\text{ds}(\mathcal{T}) = m$, $\text{RS}(\mathcal{T}') = \beta$, $\text{ds}(\mathcal{T}') = n$.

¹⁸Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Topologies for families of complete theories(with S.V.Sudoplatov)¹⁸

- We consider families \mathcal{T} of consistent first-order theories in languages $\Sigma \subseteq \Sigma(\mathcal{T})$, not necessary complete. By $F(\Sigma)$ we denote the set of all formulas in the language Σ and by $\text{Sent}(\Sigma)$ the set of all sentences in $F(\Sigma)$.
- **Proposition 7.** Any family \mathcal{T} forms a T_0 -space.
- **Proposition 8.** For any family \mathcal{T} the following conditions are equivalent:
 - (1) \mathcal{T} is a T_1 -space;
 - (2) \mathcal{T} does not contain theories T, T' with $T \subsetneq T'$.
- **Proposition 9.** A T_1 -space \mathcal{T} is Hausdorff if and only if for any distinct theories $T, T' \in \mathcal{T}$ there are sentences $\varphi \in T, \psi \in T'$ such that $\varphi \wedge \psi$ is \mathcal{T} -inconsistent.
- **Theorem 10.** For any ordinals $\alpha \leq \beta$ and natural $m, n \in \omega \setminus \{0\}$, where $m \leq n$ if $\beta = \alpha$, there is a family \mathcal{T} and its Σ' -restriction \mathcal{T}' such that $\text{RS}(\mathcal{T}) = \alpha, \text{ds}(\mathcal{T}) = m, \text{RS}(\mathcal{T}') = \beta, \text{ds}(\mathcal{T}') = n$.

¹⁸Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Topologies for families of complete theories(with S.V.Sudoplatov)¹⁸

- We consider families \mathcal{T} of consistent first-order theories in languages $\Sigma \subseteq \Sigma(\mathcal{T})$, not necessary complete. By $F(\Sigma)$ we denote the set of all formulas in the language Σ and by $\text{Sent}(\Sigma)$ the set of all sentences in $F(\Sigma)$.
- **Proposition 7.** Any family \mathcal{T} forms a T_0 -space.
- **Proposition 8.** For any family \mathcal{T} the following conditions are equivalent:
 - (1) \mathcal{T} is a T_1 -space;
 - (2) \mathcal{T} does not contain theories T, T' with $T \subsetneq T'$.
- **Proposition 9.** A T_1 -space \mathcal{T} is Hausdorff if and only if for any distinct theories $T, T' \in \mathcal{T}$ there are sentences $\varphi \in T, \psi \in T'$ such that $\varphi \wedge \psi$ is \mathcal{T} -inconsistent.
- **Theorem 10.** For any ordinals $\alpha \leq \beta$ and natural $m, n \in \omega \setminus \{0\}$, where $m \leq n$ if $\beta = \alpha$, there is a family \mathcal{T} and its Σ' -restriction \mathcal{T}' such that $\text{RS}(\mathcal{T}) = \alpha, \text{ds}(\mathcal{T}) = m, \text{RS}(\mathcal{T}') = \beta, \text{ds}(\mathcal{T}') = n$.

¹⁸Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

\bar{e} -minimal and Bs -minimal subfamilies(with S.V.Sudoplatov)¹⁹

- **Definition.** An infinite family \mathcal{T} of (possibly incomplete) theories is called *\bar{e} -minimal* if for any sentence $\varphi \in \text{Sent}(\Sigma(\mathcal{T}))$, \mathcal{T}_φ is finite or $\overline{\mathcal{T}_\varphi} = \mathcal{T} \setminus \mathcal{T}_\varphi$ is finite.
- Any boolean combination of s -definable subfamilies of \mathcal{T} is reduced to unions of sets of form $\mathcal{T}_\varphi \cap \overline{\mathcal{T}_{\psi_1}} \cap \dots \cap \overline{\mathcal{T}_{\psi_n}}$, which, in general case, can not be written shorter. These boolean combinations are called *Bs -definable* subfamilies of \mathcal{T} .
- For a family \mathcal{T} we denote by $Bs_{\mathcal{T}}$ the set of all Bs -definable subfamilies of \mathcal{T} . Clearly, $Bs_{\mathcal{T}}$ forms a boolean algebra. Moreover, we have:
- **Lemma 11.** *For any family \mathcal{T} the pair $(\mathcal{T}, Bs_{\mathcal{T}})$ is Hausdorff.*

¹⁹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

\bar{e} -minimal and Bs -minimal subfamilies(with S.V.Sudoplatov)¹⁹

- **Definition.** An infinite family \mathcal{T} of (possibly incomplete) theories is called *\bar{e} -minimal* if for any sentence $\varphi \in \text{Sent}(\Sigma(\mathcal{T}))$, \mathcal{T}_φ is finite or $\overline{\mathcal{T}_\varphi} = \mathcal{T} \setminus \mathcal{T}_\varphi$ is finite.
- Any boolean combination of s -definable subfamilies of \mathcal{T} is reduced to unions of sets of form $\mathcal{T}_\varphi \cap \overline{\mathcal{T}_{\psi_1}} \cap \dots \cap \overline{\mathcal{T}_{\psi_n}}$, which, in general case, can not be written shorter. These boolean combinations are called *Bs -definable* subfamilies of \mathcal{T} .
- For a family \mathcal{T} we denote by $Bs_{\mathcal{T}}$ the set of all Bs -definable subfamilies of \mathcal{T} . Clearly, $Bs_{\mathcal{T}}$ forms a boolean algebra. Moreover, we have:
- **Lemma 11.** *For any family \mathcal{T} the pair $(\mathcal{T}, Bs_{\mathcal{T}})$ is Hausdorff.*

¹⁹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

\bar{e} -minimal and B_S -minimal subfamilies(with S.V.Sudoplatov)¹⁹

- **Definition.** An infinite family \mathcal{T} of (possibly incomplete) theories is called *\bar{e} -minimal* if for any sentence $\varphi \in \text{Sent}(\Sigma(\mathcal{T}))$, \mathcal{T}_φ is finite or $\overline{\mathcal{T}_\varphi} = \mathcal{T} \setminus \mathcal{T}_\varphi$ is finite.
- Any boolean combination of s -definable subfamilies of \mathcal{T} is reduced to unions of sets of form $\mathcal{T}_\varphi \cap \overline{\mathcal{T}_{\psi_1}} \cap \dots \cap \overline{\mathcal{T}_{\psi_n}}$, which, in general case, can not be written shorter. These boolean combinations are called *B_S -definable* subfamilies of \mathcal{T} .
- For a family \mathcal{T} we denote by $B_{S\mathcal{T}}$ the set of all B_S -definable subfamilies of \mathcal{T} . Clearly, $B_{S\mathcal{T}}$ forms a boolean algebra. Moreover, we have:
- **Lemma 11.** *For any family \mathcal{T} the pair $(\mathcal{T}, B_{S\mathcal{T}})$ is Hausdorff.*

¹⁹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

\bar{e} -minimal and B_S -minimal subfamilies(with S.V.Sudoplatov)¹⁹

- **Definition.** An infinite family \mathcal{T} of (possibly incomplete) theories is called *\bar{e} -minimal* if for any sentence $\varphi \in \text{Sent}(\Sigma(\mathcal{T}))$, \mathcal{T}_φ is finite or $\overline{\mathcal{T}_\varphi} = \mathcal{T} \setminus \mathcal{T}_\varphi$ is finite.
- Any boolean combination of s -definable subfamilies of \mathcal{T} is reduced to unions of sets of form $\mathcal{T}_\varphi \cap \overline{\mathcal{T}_{\psi_1}} \cap \dots \cap \overline{\mathcal{T}_{\psi_n}}$, which, in general case, can not be written shorter. These boolean combinations are called *B_S -definable* subfamilies of \mathcal{T} .
- For a family \mathcal{T} we denote by $B_{S\mathcal{T}}$ the set of all B_S -definable subfamilies of \mathcal{T} . Clearly, $B_{S\mathcal{T}}$ forms a boolean algebra. Moreover, we have:
- **Lemma 11.** *For any family \mathcal{T} the pair $(\mathcal{T}, B_{S\mathcal{T}})$ is Hausdorff.*

¹⁹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks for families of incomplete theories(with S.V.Sudoplatov)²⁰

- **Definition.** For the empty family \mathcal{T} we put the rank $\overline{\text{RS}}(\mathcal{T}) = -1$, and for finite nonempty families \mathcal{T} we put $\overline{\text{RS}}(\mathcal{T}) = 0$.
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if there are pairwise disjoint Bs-definable subfamilies \mathcal{T}_n of \mathcal{T} , $n \in \omega$, such that $\overline{\text{RS}}(\mathcal{T}_n) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal then $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\overline{\text{RS}}(\mathcal{T}) = \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ and $\overline{\text{RS}}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ for any ordinal α , we put $\overline{\text{RS}}(\mathcal{T}) = \infty$.

²⁰Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

<https://doi.org/10.1007/s10469-021-09620-4>

Ranks for families of incomplete theories(with S.V.Sudoplatov)²⁰

- **Definition.** For the empty family \mathcal{T} we put the rank $\overline{\text{RS}}(\mathcal{T}) = -1$, and for finite nonempty families \mathcal{T} we put $\overline{\text{RS}}(\mathcal{T}) = 0$.
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if there are pairwise disjoint Bs-definable subfamilies \mathcal{T}_n of \mathcal{T} , $n \in \omega$, such that $\overline{\text{RS}}(\mathcal{T}_n) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal then $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\overline{\text{RS}}(\mathcal{T}) = \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ and $\overline{\text{RS}}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ for any ordinal α , we put $\overline{\text{RS}}(\mathcal{T}) = \infty$.

²⁰Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks for families of incomplete theories(with S.V.Sudoplatov)²⁰

- **Definition.** For the empty family \mathcal{T} we put the rank $\overline{\text{RS}}(\mathcal{T}) = -1$, and for finite nonempty families \mathcal{T} we put $\overline{\text{RS}}(\mathcal{T}) = 0$.
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if there are pairwise disjoint Bs-definable subfamilies \mathcal{T}_n of \mathcal{T} , $n \in \omega$, such that $\overline{\text{RS}}(\mathcal{T}_n) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal then $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\overline{\text{RS}}(\mathcal{T}) = \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ and $\overline{\text{RS}}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ for any ordinal α , we put $\overline{\text{RS}}(\mathcal{T}) = \infty$.

²⁰Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks for families of incomplete theories(with S.V.Sudoplatov)²⁰

- **Definition.** For the empty family \mathcal{T} we put the rank $\overline{\text{RS}}(\mathcal{T}) = -1$, and for finite nonempty families \mathcal{T} we put $\overline{\text{RS}}(\mathcal{T}) = 0$.
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if there are pairwise disjoint Bs-definable subfamilies \mathcal{T}_n of \mathcal{T} , $n \in \omega$, such that $\overline{\text{RS}}(\mathcal{T}_n) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal then $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\overline{\text{RS}}(\mathcal{T}) = \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ and $\overline{\text{RS}}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ for any ordinal α , we put $\overline{\text{RS}}(\mathcal{T}) = \infty$.

²⁰Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks for families of incomplete theories(with S.V.Sudoplatov)²⁰

- **Definition.** For the empty family \mathcal{T} we put the rank $\overline{\text{RS}}(\mathcal{T}) = -1$, and for finite nonempty families \mathcal{T} we put $\overline{\text{RS}}(\mathcal{T}) = 0$.
- For a family \mathcal{T} and an ordinal $\alpha = \beta + 1$ we put $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if there are pairwise disjoint Bs-definable subfamilies \mathcal{T}_n of \mathcal{T} , $n \in \omega$, such that $\overline{\text{RS}}(\mathcal{T}_n) \geq \beta$, $n \in \omega$.
- If α is a limit ordinal then $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \beta$ for any $\beta < \alpha$.
- We set $\overline{\text{RS}}(\mathcal{T}) = \alpha$ if $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ and $\overline{\text{RS}}(\mathcal{T}) \not\geq \alpha + 1$.
- If $\overline{\text{RS}}(\mathcal{T}) \geq \alpha$ for any ordinal α , we put $\overline{\text{RS}}(\mathcal{T}) = \infty$.

²⁰Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks and degrees for families of incomplete theories(with S.V.Sudoplatov)²¹

- **Definition.** A family \mathcal{T} is called \bar{e} -totally transcendental if $\overline{\text{RS}}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is \bar{e} -totally transcendental, with $\overline{\text{RS}}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\overline{\text{ds}}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise disjoint B_s-definable subfamilies \mathcal{T}_i such that $\overline{\text{RS}}(\mathcal{T}_i) = \alpha$.
- By the definition, if $\overline{\text{RS}}(\mathcal{T}) = \alpha$, for an ordinal α , then $\overline{\text{ds}}(\mathcal{T}) \in \omega \setminus \{0\}$.
- **Theorem 12.** For any ordinal α and natural $n \neq 0$ there is a family \mathcal{T} of positive theories such that $(\overline{\text{RS}}(\mathcal{T}), \overline{\text{ds}}(\mathcal{T})) = (\alpha, n)$.

²¹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks and degrees for families of incomplete theories(with S.V.Sudoplatov)²¹

- **Definition.** A family \mathcal{T} is called \bar{e} -totally transcendental if $\overline{\text{RS}}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is \bar{e} -totally transcendental, with $\overline{\text{RS}}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\overline{\text{ds}}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise disjoint B_S-definable subfamilies \mathcal{T}_i such that $\overline{\text{RS}}(\mathcal{T}_i) = \alpha$.
- By the definition, if $\overline{\text{RS}}(\mathcal{T}) = \alpha$, for an ordinal α , then $\overline{\text{ds}}(\mathcal{T}) \in \omega \setminus \{0\}$.
- **Theorem 12.** For any ordinal α and natural $n \neq 0$ there is a family \mathcal{T} of positive theories such that $(\overline{\text{RS}}(\mathcal{T}), \overline{\text{ds}}(\mathcal{T})) = (\alpha, n)$.

²¹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks and degrees for families of incomplete theories(with S.V.Sudoplatov)²¹

- **Definition.** A family \mathcal{T} is called \bar{e} -totally transcendental if $\overline{\text{RS}}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is \bar{e} -totally transcendental, with $\overline{\text{RS}}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\overline{\text{ds}}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise disjoint Bs -definable subfamilies \mathcal{T}_i such that $\overline{\text{RS}}(\mathcal{T}_i) = \alpha$.
- By the definition, if $\overline{\text{RS}}(\mathcal{T}) = \alpha$, for an ordinal α , then $\overline{\text{ds}}(\mathcal{T}) \in \omega \setminus \{0\}$.
- **Theorem 12.** *For any ordinal α and natural $n \neq 0$ there is a family \mathcal{T} of positive theories such that $(\overline{\text{RS}}(\mathcal{T}), \overline{\text{ds}}(\mathcal{T})) = (\alpha, n)$.*

²¹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Ranks and degrees for families of incomplete theories(with S.V.Sudoplatov)²¹

- **Definition.** A family \mathcal{T} is called \bar{e} -totally transcendental if $\overline{\text{RS}}(\mathcal{T})$ is an ordinal.
- If \mathcal{T} is \bar{e} -totally transcendental, with $\overline{\text{RS}}(\mathcal{T}) = \alpha \geq 0$, we define the *degree* $\overline{\text{ds}}(\mathcal{T})$ of \mathcal{T} as the maximal number of pairwise disjoint Bs -definable subfamilies \mathcal{T}_i such that $\overline{\text{RS}}(\mathcal{T}_i) = \alpha$.
- By the definition, if $\overline{\text{RS}}(\mathcal{T}) = \alpha$, for an ordinal α , then $\overline{\text{ds}}(\mathcal{T}) \in \omega \setminus \{0\}$.
- **Theorem 12.** For any ordinal α and natural $n \neq 0$ there is a family \mathcal{T} of positive theories such that $(\overline{\text{RS}}(\mathcal{T}), \overline{\text{ds}}(\mathcal{T})) = (\alpha, n)$.

²¹Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²²

- An intersections, possibly infinite, of B_S -definable subfamilies of \mathcal{T} are called *B_S -diagram definable*, or *Bd-definable*.
- We extend the algebra $\mathcal{B}_{B_S}(\mathcal{T})$ till the algebra $\mathcal{B}_{Bd}(\mathcal{T})$ of all Bd-definable subfamilies of \mathcal{T} . Clearly, the algebra $\mathcal{B}_{Bd}(\mathcal{T})$ preserves the operations \cap and \cup , whereas complements are defined only for B_S -definable sets. Therefore $\mathcal{B}_{Bd}(\mathcal{T})$ contains the operations \cap and \cup , being a partial algebra with respect to $\bar{}$.
- Thus, $\mathcal{B}_{Bd}(\mathcal{T})$ forms a distributive lattice, partially, for B_S -definable sets, with complements.

²²Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

<https://doi.org/10.1007/s10469-021-09620-4>

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²²

- An intersections, possibly infinite, of B_S -definable subfamilies of \mathcal{T} are called B_S -*diagram definable*, or B_d -*definable*.
- We extend the algebra $\mathcal{B}_{B_S}(\mathcal{T})$ till the algebra $\mathcal{B}_{B_d}(\mathcal{T})$ of all B_d -definable subfamilies of \mathcal{T} . Clearly, the algebra $\mathcal{B}_{B_d}(\mathcal{T})$ preserves the operations \cap and \cup , whereas complements are defined only for B_S -definable sets. Therefore $\mathcal{B}_{B_d}(\mathcal{T})$ contains the operations \cap and \cup , being a partial algebra with respect to $\bar{}$.
- Thus, $\mathcal{B}_{B_d}(\mathcal{T})$ forms a distributive lattice, partially, for B_S -definable sets, with complements.

²²Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²²

- An intersections, possibly infinite, of B_S -definable subfamilies of \mathcal{T} are called *B_S -diagram definable*, or *B_d -definable*.
- We extend the algebra $\mathcal{B}_{B_S}(\mathcal{T})$ till the algebra $\mathcal{B}_{B_d}(\mathcal{T})$ of all B_d -definable subfamilies of \mathcal{T} . Clearly, the algebra $\mathcal{B}_{B_d}(\mathcal{T})$ preserves the operations \cap and \cup , whereas complements are defined only for B_S -definable sets. Therefore $\mathcal{B}_{B_d}(\mathcal{T})$ contains the operations \cap and \cup , being a partial algebra with respect to $\bar{}$.
- Thus, $\mathcal{B}_{B_d}(\mathcal{T})$ forms a distributive lattice, partially, for B_S -definable sets, with complements.

²²Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²³

- **Theorem 13.** *For any nonempty B_S -closed special family \mathcal{T} , an ordinal $\alpha \geq 1$, and $n \in \omega \setminus \{0\}$, the following conditions are equivalent: (1) $\overline{RS}(\mathcal{T}) = \alpha$ and $\overline{ds}(\mathcal{T}) = n$;*
- (2) *the Boolean algebra $\mathcal{B}_{B_S}(\mathcal{T})$ is isomorphic to a direct product of n Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$ each of which is generated by elements of ranks $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
- (3) *the algebra $\mathcal{B}_{Bd}(\mathcal{T})$ consists of infinitely many atomic elements of each rank $\beta < \alpha$, for $\beta \geq 0$, and exactly n atomic elements of rank α , these n atomic elements correspond non-principal ultrafilters with respect to elements of ranks $< \alpha$.*

²³Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²³

- **Theorem 13.** *For any nonempty B_S -closed special family \mathcal{T} , an ordinal $\alpha \geq 1$, and $n \in \omega \setminus \{0\}$, the following conditions are equivalent: (1) $\overline{RS}(\mathcal{T}) = \alpha$ and $\overline{ds}(\mathcal{T}) = n$;*
- (2) *the Boolean algebra $\mathcal{B}_{B_S}(\mathcal{T})$ is isomorphic to a direct product of n Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$ each of which is generated by elements of ranks $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
- (3) *the algebra $\mathcal{B}_{Bd}(\mathcal{T})$ consists of infinitely many atomic elements of each rank $\beta < \alpha$, for $\beta \geq 0$, and exactly n atomic elements of rank α , these n atomic elements correspond non-principal ultrafilters with respect to elements of ranks $< \alpha$.*

²³Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

Algebras for B_S -definable subfamilies of families of theories (with S.V.Sudoplatov)²³

- **Theorem 13.** *For any nonempty B_S -closed special family \mathcal{T} , an ordinal $\alpha \geq 1$, and $n \in \omega \setminus \{0\}$, the following conditions are equivalent: (1) $\overline{RS}(\mathcal{T}) = \alpha$ and $\overline{ds}(\mathcal{T}) = n$;*
- (2) *the Boolean algebra $\mathcal{B}_{B_S}(\mathcal{T})$ is isomorphic to a direct product of n Boolean algebras $\mathcal{B}_1, \dots, \mathcal{B}_n$ each of which is generated by elements of ranks $< \alpha$ such that each \mathcal{B}_n contains infinitely many elements of each rank $\beta < \alpha$;*
- (3) *the algebra $\mathcal{B}_{Bd}(\mathcal{T})$ consists of infinitely many atomic elements of each rank $\beta < \alpha$, for $\beta \geq 0$, and exactly n atomic elements of rank α , these n atomic elements correspond non-principal ultrafilters with respect to elements of ranks $< \alpha$.*

²³Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. I. Algebra and Logic 59:6, 437–455 (2021).

- **Definition.** For a family \mathcal{T} of theories in a language Σ and a theory T we put $T \in \text{Cl}_1(\mathcal{T})$ if $T \in \mathcal{T}$, or T is nonempty and

$$T = \{\varphi \in \text{Sent}(\Sigma) \mid (\mathcal{T}')_\varphi \text{ is infinite}\}$$

for some $\mathcal{T}' \subseteq \mathcal{T}$.

- If \mathcal{T}' is fixed then we say that T belongs to the Cl_1 -closure of \mathcal{T} with respect to \mathcal{T}' , and T is an accumulation point of \mathcal{T} with respect to \mathcal{T}' .
- A family \mathcal{T} is called Cl_1 -closed or simply closed if $\mathcal{T} = \text{Cl}_1(\mathcal{T})$. We assume that any $T \in \text{Cl}_1(\mathcal{T}) \setminus \mathcal{T}$ is nonempty, since otherwise the empty theory always belongs to $\text{Cl}_1(\mathcal{T})$, using $\mathcal{T}' = \emptyset$.

²⁴Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

- **Definition.** For a family \mathcal{T} of theories in a language Σ and a theory T we put $T \in \text{Cl}_1(\mathcal{T})$ if $T \in \mathcal{T}$, or T is nonempty and

$$T = \{\varphi \in \text{Sent}(\Sigma) \mid (\mathcal{T}')_\varphi \text{ is infinite}\}$$

for some $\mathcal{T}' \subseteq \mathcal{T}$.

- If \mathcal{T}' is fixed then we say that T *belongs to the Cl_1 -closure of \mathcal{T} with respect to \mathcal{T}'* , and T is an *accumulation point* of \mathcal{T} with respect to \mathcal{T}' .
- A family \mathcal{T} is called *Cl_1 -closed* or simply *closed* if $\mathcal{T} = \text{Cl}_1(\mathcal{T})$. We assume that any $T \in \text{Cl}_1(\mathcal{T}) \setminus \mathcal{T}$ is nonempty, since otherwise the empty theory always belongs to $\text{Cl}_1(\mathcal{T})$, using $\mathcal{T}' = \emptyset$.

²⁴Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

- **Definition.** For a family \mathcal{T} of theories in a language Σ and a theory T we put $T \in \text{Cl}_1(\mathcal{T})$ if $T \in \mathcal{T}$, or T is nonempty and

$$T = \{\varphi \in \text{Sent}(\Sigma) \mid (\mathcal{T}')_\varphi \text{ is infinite}\}$$

for some $\mathcal{T}' \subseteq \mathcal{T}$.

- If \mathcal{T}' is fixed then we say that T *belongs to the Cl_1 -closure of \mathcal{T} with respect to \mathcal{T}'* , and T is an *accumulation point* of \mathcal{T} with respect to \mathcal{T}' .
- A family \mathcal{T} is called *Cl_1 -closed* or simply *closed* if $\mathcal{T} = \text{Cl}_1(\mathcal{T})$. We assume that any $T \in \text{Cl}_1(\mathcal{T}) \setminus \mathcal{T}$ is nonempty, since otherwise the empty theory always belongs to $\text{Cl}_1(\mathcal{T})$, using $\mathcal{T}' = \emptyset$.

²⁴Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

<https://doi.org/10.33048/alglog.2021.60.104>

- **Theorem 14.** *For any family \mathcal{T} of complete theories in at most countable language Σ , $\text{Cl}_1(\mathcal{T}) = \text{Cl}_E(\mathcal{T})$.*
- **Theorem 15.** *For any family \mathcal{T} in at most countable language Σ the closure operator Cl_1 is transitive.*
- **Theorem 16.** *For any family \mathcal{T} of theories in a countable language the following conditions are equivalent:*
 - (1) $|\text{Cl}_1(\mathcal{T})| = 2^\omega$;
 - (2) $e_1\text{-Sp}(\mathcal{T}) = 2^\omega$;
 - (3) $\overline{RS}(\mathcal{T}) = \infty$.

²⁵Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

- **Theorem 14.** *For any family \mathcal{T} of complete theories in at most countable language Σ , $\text{Cl}_1(\mathcal{T}) = \text{Cl}_E(\mathcal{T})$.*
- **Theorem 15.** *For any family \mathcal{T} in at most countable language Σ the closure operator Cl_1 is transitive.*
- **Theorem 16.** *For any family \mathcal{T} of theories in a countable language the following conditions are equivalent:*
 - (1) $|\text{Cl}_1(\mathcal{T})| = 2^\omega$;
 - (2) $e_1\text{-Sp}(\mathcal{T}) = 2^\omega$;
 - (3) $\overline{RS}(\mathcal{T}) = \infty$.

²⁵Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

- **Theorem 14.** *For any family \mathcal{T} of complete theories in at most countable language Σ , $\text{Cl}_1(\mathcal{T}) = \text{Cl}_E(\mathcal{T})$.*
- **Theorem 15.** *For any family \mathcal{T} in at most countable language Σ the closure operator Cl_1 is transitive.*
- **Theorem 16.** *For any family \mathcal{T} of theories in a countable language the following conditions are equivalent:*
 - (1) $|\text{Cl}_1(\mathcal{T})| = 2^\omega$;
 - (2) $e_1\text{-Sp}(\mathcal{T}) = 2^\omega$;
 - (3) $\overline{RS}(\mathcal{T}) = \infty$.

²⁵Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

Closures for linearly ordered families of theories(with S.V.Sudoplatov)²⁶

- **Theorem 17.** *For any linearly \subseteq -ordered family \mathcal{T} , $\text{Cl}_1(\mathcal{T})$ consists of unions for subfamilies of \mathcal{T} , and of intersections for countable subfamilies of \mathcal{T} ordered by the type ω^* .*
- **Theorem 18.** *For any linearly \subseteq -ordered family \mathcal{T} , $\text{Cl}_1(\text{Cl}_1(\mathcal{T})) = \text{Cl}_1(\mathcal{T})$.*

²⁶Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

<https://doi.org/10.33048/alglog.2021.60.104>

Closures for linearly ordered families of theories(with S.V.Sudoplatov)²⁶

- **Theorem 17.** *For any linearly \subseteq -ordered family \mathcal{T} , $\text{Cl}_1(\mathcal{T})$ consists of unions for subfamilies of \mathcal{T} , and of intersections for countable subfamilies of \mathcal{T} ordered by the type ω^* .*
- **Theorem 18.** *For any linearly \subseteq -ordered family \mathcal{T} , $\text{Cl}_1(\text{Cl}_1(\mathcal{T})) = \text{Cl}_1(\mathcal{T})$.*

²⁶Markhabatov, N.D., Sudoplatov, S.V. Topologies, Ranks, and Closures for Families of Theories. II. Algebra and Logic 60:1, 57-80 (2021).

<https://doi.org/10.33048/alglog.2021.60.104>

The family of permutation theories²⁷

- Let a language Σ consist of the permutation f . Denote by \mathcal{T}_Σ family of all permutation theories of language Σ .
- The following theorem shows that there is a family of permutations of theory, having a countable rank.
- **Theorem 19.** *For any countable ordinal α and natural $k \geq 1$ there exists a family $\mathcal{T} \subseteq \mathcal{T}_\Sigma$, such that $RS(\mathcal{T}) = \alpha$ and $ds(\mathcal{T}) = k$.*
- **Theorem 20.** *Any theory T permutation on an infinite set is pseudofinite.*

²⁷Markhabatov, N. D. Ranks for families of permutation theories / N. D. Markhabatov // Известия Иркутского государственного университета. Серия: Математика. — 2019. — Т. 28. — С. 85—94.

The family of permutation theories²⁷

- Let a language Σ consist of the permutation f . Denote by \mathcal{T}_Σ family of all permutation theories of language Σ .
- The following theorem shows that there is a family of permutations of theory, having a countable rank.
- *Theorem 19. For any countable ordinal α and natural $k \geq 1$ there exists a family $\mathcal{T} \subseteq \mathcal{T}_\Sigma$, such that $RS(\mathcal{T}) = \alpha$ and $ds(\mathcal{T}) = k$.*
- *Theorem 20. Any theory T permutation on an infinite set is pseudofinite.*

²⁷Markhabatov, N. D. Ranks for families of permutation theories / N. D. Markhabatov // Известия Иркутского государственного университета. Серия: Математика. — 2019. — Т. 28. — С. 85—94.

The family of permutation theories²⁷

- Let a language Σ consist of the permutation f . Denote by \mathcal{T}_Σ family of all permutation theories of language Σ .
- The following theorem shows that there is a family of permutations of theory, having a countable rank.
- **Theorem 19.** *For any countable ordinal α and natural $k \geq 1$ there exists a family $\mathcal{T} \subseteq \mathcal{T}_\Sigma$, such that $RS(\mathcal{T}) = \alpha$ and $ds(\mathcal{T}) = k$.*
- **Theorem 20.** *Any theory T permutation on an infinite set is pseudofinite.*

²⁷Markhabatov, N. D. Ranks for families of permutation theories / N. D. Markhabatov // Известия Иркутского государственного университета. Серия: Математика. — 2019. — Т. 28. — С. 85—94.

The family of permutation theories²⁷

- Let a language Σ consist of the permutation f . Denote by \mathcal{T}_Σ family of all permutation theories of language Σ .
- The following theorem shows that there is a family of permutations of theory, having a countable rank.
- **Theorem 19.** *For any countable ordinal α and natural $k \geq 1$ there exists a family $\mathcal{T} \subseteq \mathcal{T}_\Sigma$, such that $RS(\mathcal{T}) = \alpha$ and $ds(\mathcal{T}) = k$.*
- **Theorem 20.** *Any theory T permutation on an infinite set is pseudofinite.*

²⁷Markhabatov, N. D. Ranks for families of permutation theories / N. D. Markhabatov // Известия Иркутского государственного университета. Серия: Математика. — 2019. — Т. 28. — С. 85—94.

- Let T be a complete first-order theory of locally free algebras of signature $\Sigma = \{f_1^{(n_1)}, \dots, f_r^{(n_r)}, \dots; c_1, \dots, c_k, \dots, \}$.
- *Theorem 21. An infinite locally free algebra A is pseudofinite if and only if the signature contains no more than one unary symbol and does not contain symbols arity $n > 1$.*
- *Corollary 22. A locally free groupoid is not pseudofinite.*

²⁸Markhabatov, N. D. Pseudofiniteness of locally free algebras / N. D. Markhabatov // Algebra and model theory 12 : coll. of papers. — Novosibirsk : NSTU publ. — 2019. — T. 12. — C. 41—46.

- Let T be a complete first-order theory of locally free algebras of signature $\Sigma = \{f_1^{(n_1)}, \dots, f_r^{(n_r)}, \dots; c_1, \dots, c_k, \dots, \}$.
- *Theorem 21. An infinite locally free algebra A is pseudofinite if and only if the signature contains no more than one unary symbol and does not contain symbols arity $n > 1$.*
- *Corollary 22. A locally free groupoid is not pseudofinite.*

²⁸Markhabatov, N. D. Pseudofiniteness of locally free algebras / N. D. Markhabatov // Algebra and model theory 12 : coll. of papers. — Novosibirsk : NSTU publ. — 2019. — T. 12. — C. 41—46.

- Let T be a complete first-order theory of locally free algebras of signature $\Sigma = \{f_1^{(n_1)}, \dots, f_r^{(n_r)}, \dots; c_1, \dots, c_k, \dots, \}$.
- **Theorem 21.** *An infinite locally free algebra A is pseudofinite if and only if the signature contains no more than one unary symbol and does not contain symbols arity $n > 1$.*
- **Corollary 22.** *A locally free groupoid is not pseudofinite.*

²⁸Markhabatov, N. D. Pseudofiniteness of locally free algebras / N. D. Markhabatov // Algebra and model theory 12 : coll. of papers. — Novosibirsk : NSTU publ. — 2019. — T. 12. — C. 41—46.

- Let T be a complete first-order theory of locally free algebras of signature $\Sigma = \{f_1^{(n_1)}, \dots, f_r^{(n_r)}, \dots; c_1, \dots, c_k, \dots, \}$.
- **Theorem 21.** *An infinite locally free algebra A is pseudofinite if and only if the signature contains no more than one unary symbol and does not contain symbols arity $n > 1$.*
- **Corollary 22.** *A locally free groupoid is not pseudofinite.*

²⁸Markhabatov, N. D. Pseudofiniteness of locally free algebras / N. D. Markhabatov // Algebra and model theory 12 : coll. of papers. – Novosibirsk : NSTU publ. — 2019. — T. 12. — C. 41—46.

Thank you!