

CANCELLABLE ELEMENTS OF THE LATTICE OF MONOID VARIETIES

S.V. Gusev, E.W.H. Lee

Ural Federal University, Yekaterinburg, Russia
Nova Southeastern University, Fort Lauderdale, USA

14th International Summer School-Conference
"Problems Allied to Model Theory and Universal Algebra"
Erlagol, June 27, 2021

A non-empty class of universal algebras of the same type is called a *variety* if it is closed under subalgebras, homomorphic images and direct products.

A class of universal algebras is a variety if and only if it is axiomatized by a set of identities (Birkhoff's theorem).

The collection of all varieties forms a lattice under class-theoretical inclusion.

L.N. Shevrin, B.M. Vernikov and M.V. Volkov, Lattices of semigroup varieties, Russ. Math. Izv. VUZ, 53, No. 3 (2009), 1–28.

A non-empty class of universal algebras of the same type is called a *variety* if it is closed under subalgebras, homomorphic images and direct products.

A class of universal algebras is a variety if and only if it is axiomatized by a set of identities (Birkhoff's theorem).

The collection of all varieties forms a lattice under class-theoretical inclusion.

L.N. Shevrin, B.M. Vernikov and M.V. Volkov, Lattices of semigroup varieties, Russ. Math. Izv. VUZ, 53, No. 3 (2009), 1–28.

A non-empty class of universal algebras of the same type is called a *variety* if it is closed under subalgebras, homomorphic images and direct products.

A class of universal algebras is a variety if and only if it is axiomatized by a set of identities (Birkhoff's theorem).

The collection of all varieties forms a lattice under class-theoretical inclusion.

L.N. Shevrin, B.M. Vernikov and M.V. Volkov, Lattices of semigroup varieties, Russ. Math. Izv. VUZ, 53, No. 3 (2009), 1–28.

A non-empty class of universal algebras of the same type is called a *variety* if it is closed under subalgebras, homomorphic images and direct products.

A class of universal algebras is a variety if and only if it is axiomatized by a set of identities (Birkhoff's theorem).

The collection of all varieties forms a lattice under class-theoretical inclusion.

L.N. Shevrin, B.M. Vernikov and M.V. Volkov, Lattices of semigroup varieties, Russ. Math. Izv. VUZ, 53, No. 3 (2009), 1–28.

Let $I(x_1, x_2, \dots, x_n)$ be a lattice identity or a lattice quasi-identity.
An element a of a lattice L is an I -element if

$$(\forall x_2, x_3, \dots, x_n \in L) \quad I(a, x_2, x_3, \dots, x_n).$$

An element a of a lattice L is called *neutral* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge (x \vee y) \wedge (y \vee a) = (a \wedge x) \vee (x \wedge y) \vee (y \wedge a).$$

$[a] = \{x \in L \mid x \leq a\}$ is the principal ideal generated by a .

$[a] = \{x \in L \mid x \geq a\}$ is the principal filter generated by a .

An element a of L is neutral if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an embedding from L into $[a] \times [a]$.

Let $I(x_1, x_2, \dots, x_n)$ be a lattice identity or a lattice quasi-identity.
An element a of a lattice L is an I -element if

$$(\forall x_2, x_3, \dots, x_n \in L) \quad I(a, x_2, x_3, \dots, x_n).$$

An element a of a lattice L is called *neutral* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge (x \vee y) \wedge (y \vee a) = (a \wedge x) \vee (x \wedge y) \vee (y \wedge a).$$

$[a] = \{x \in L \mid x \leq a\}$ is the principal ideal generated by a .

$[a] = \{x \in L \mid x \geq a\}$ is the principal filter generated by a .

An element a of L is neutral if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an embedding from L into $[a] \times [a]$.

Let $I(x_1, x_2, \dots, x_n)$ be a lattice identity or a lattice quasi-identity.
An element a of a lattice L is an I -element if

$$(\forall x_2, x_3, \dots, x_n \in L) \quad I(a, x_2, x_3, \dots, x_n).$$

An element a of a lattice L is called *neutral* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge (x \vee y) \wedge (y \vee a) = (a \wedge x) \vee (x \wedge y) \vee (y \wedge a).$$

$[a] = \{x \in L \mid x \leq a\}$ is the principal ideal generated by a .

$[a] = \{x \in L \mid x \geq a\}$ is the principal filter generated by a .

An element a of L is neutral if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an embedding from L into $[a] \times [a]$.

An element a is called *standard* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge y = (a \wedge y) \vee (x \wedge y).$$

Costandard elements are dual to standard ones.

G. Grätzer, Lattice Theory: Foundation, Springer Basel AG, 2011.

An element a of a lattice L is called *cancellable* if

$$(\forall x, y \in L) \quad a \vee x = a \vee y \ \& \ a \wedge x = a \wedge y \longrightarrow x = y.$$

An element a of L is cancellable if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an injection from L into $[a] \times [a]$.

A neutral element is both standard and costandard.

A standard or costandard element is cancellable.

An element a is called *standard* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge y = (a \wedge y) \vee (x \wedge y).$$

Costandard elements are dual to standard ones.

G. Grätzer, Lattice Theory: Foundation, Springer Basel AG, 2011.

An element a of a lattice L is called *cancellable* if

$$(\forall x, y \in L) \quad a \vee x = a \vee y \ \& \ a \wedge x = a \wedge y \longrightarrow x = y.$$

An element a of L is cancellable if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an injection from L into $[a] \times [a]$.

A neutral element is both standard and costandard.

A standard or costandard element is cancellable.

An element a is called *standard* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge y = (a \wedge y) \vee (x \wedge y).$$

Costandard elements are dual to standard ones.

G. Grätzer, Lattice Theory: Foundation, Springer Basel AG, 2011.

An element a of a lattice L is called *cancellable* if

$$(\forall x, y \in L) \quad a \vee x = a \vee y \ \& \ a \wedge x = a \wedge y \longrightarrow x = y.$$

An element a of L is cancellable if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an injection from L into $[a] \times [a]$.

A neutral element is both standard and costandard.

A standard or costandard element is cancellable.

An element a is called *standard* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge y = (a \wedge y) \vee (x \wedge y).$$

Costandard elements are dual to standard ones.

G. Grätzer, Lattice Theory: Foundation, Springer Basel AG, 2011.

An element a of a lattice L is called *cancellable* if

$$(\forall x, y \in L) \quad a \vee x = a \vee y \ \& \ a \wedge x = a \wedge y \longrightarrow x = y.$$

An element a of L is cancellable if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an injection from L into $[a] \times [a]$.

A neutral element is both standard and costandard.

A standard or costandard element is cancellable.

An element a is called *standard* if

$$(\forall x, y \in L) \quad (a \vee x) \wedge y = (a \wedge y) \vee (x \wedge y).$$

Costandard elements are dual to standard ones.

G. Grätzer, Lattice Theory: Foundation, Springer Basel AG, 2011.

An element a of a lattice L is called *cancellable* if

$$(\forall x, y \in L) \quad a \vee x = a \vee y \ \& \ a \wedge x = a \wedge y \longrightarrow x = y.$$

An element a of L is cancellable if and only if $\varphi_a: x \mapsto (x \wedge a, x \vee a)$ is an injection from L into $[a] \times [a]$.

A neutral element is both standard and costandard.

A standard or costandard element is cancellable.

B.M. Vernikov, Special elements in lattices of semigroup varieties, *Acta Sci. Math.* (Szeged), 81 (2015), 79–109.

V. Yu. Shaprynskiĭ, D.V. Skokov, B.M. Vernikov, Cancellable elements of the lattices of varieties of semigroups and epigroups, *Comm. Algebra*, 47 (2019), 4697–4712.

S.V. Gusev, Special elements of the lattice of monoid varieties, *Algebra Universalis*, 79 (2018), Article 29, 1–12.

S.V. Gusev, Standard elements of the lattice of monoid varieties, *Algebra and Logic*, 59, No. 6 (2020), 415–422.

B.M. Vernikov, Special elements in lattices of semigroup varieties, *Acta Sci. Math.* (Szeged), 81 (2015), 79–109.

V. Yu. Shaprynskiĭ, D.V. Skokov, B.M. Vernikov, Cancellable elements of the lattices of varieties of semigroups and epigroups, *Comm. Algebra*, 47 (2019), 4697–4712.

S.V. Gusev, Special elements of the lattice of monoid varieties, *Algebra Universalis*, 79 (2018), Article 29, 1–12.

S.V. Gusev, Standard elements of the lattice of monoid varieties, *Algebra and Logic*, 59, No. 6 (2020), 415–422.

B.M. Vernikov, Special elements in lattices of semigroup varieties, *Acta Sci. Math. (Szeged)*, 81 (2015), 79–109.

V. Yu. Shaprynskiĭ, D.V. Skokov, B.M. Vernikov, Cancellable elements of the lattices of varieties of semigroups and epigroups, *Comm. Algebra*, 47 (2019), 4697–4712.

S.V. Gusev, Special elements of the lattice of monoid varieties, *Algebra Universalis*, 79 (2018), Article 29, 1–12.

S.V. Gusev, Standard elements of the lattice of monoid varieties, *Algebra and Logic*, 59, No. 6 (2020), 415–422.

\mathbf{MON} denote the lattice of monoid varieties.

\mathbf{T} is the trivial variety of monoids.

\mathbf{SL} is the variety of all semilattice monoids.

\mathbf{MON} is the variety of all monoids.

Theorem (Gusev, 2018, 2020)

For a monoid variety V , the following are equivalent:

- (i) V is a neutral element of the lattice \mathbf{MON} ;
- (ii) V is a standard element of the lattice \mathbf{MON} ;
- (iii) V is one of the varieties \mathbf{T} , \mathbf{SL} or \mathbf{MON} .

The set of all neutral elements of the lattice of semigroup varieties is 5-element (Volkov, 2005).

The set of all standard elements of the lattice of semigroup varieties is countably infinite. (Shaprynskiĭ and Vernikov, 2010).

\mathbf{MON} denote the lattice of monoid varieties.

\mathbf{T} is the trivial variety of monoids.

\mathbf{SL} is the variety of all semilattice monoids.

\mathbf{MON} is the variety of all monoids.

Theorem (Gusev, 2018, 2020)

For a monoid variety V , the following are equivalent:

- (i) V is a neutral element of the lattice \mathbf{MON} ;
- (ii) V is a standard element of the lattice \mathbf{MON} ;
- (iii) V is one of the varieties \mathbf{T} , \mathbf{SL} or \mathbf{MON} .

The set of all neutral elements of the lattice of semigroup varieties is 5-element (Volkov, 2005).

The set of all standard elements of the lattice of semigroup varieties is countably infinite. (Shaprynskiĭ and Vernikov, 2010).

$$C = \text{var}\{x^2 \approx x^3, xy \approx yx\}.$$

Theorem (Gusev, 2018)

The varieties T , SL , C and MON are only costandard elements of MON .

An element of the lattice of semigroup varieties is costandard if and only if it is neutral (Vernikov, 2011).

$$D = \text{var}\{x^2 \approx x^3, x^2y \approx xyx \approx yx^2\}.$$

Theorem


The varieties T, SL, C, D and MON are only cancellable elements of MON.

The set of all cancellable elements of the lattice of semigroup varieties is countably infinite (Shaprynskiĭ, Skokov and Vernikov, 2019).

Corollary

The class of all cancellable elements of \mathbf{MON} forms a sublattice in \mathbf{MON} .

In general, the set of cancellable elements in a lattice need not form a sublattice.



An element a of a lattice L is called *modular* if

$$(\forall x, y \in L) \quad x \leq y \longrightarrow (a \vee x) \wedge y = (a \wedge y) \vee x.$$

Necessary condition of the modularity

Let V be a proper monoid variety that is a modular element of the lattice **MON**. Then V satisfies the identities

$$x^2 \approx x^3 \text{ and } x^2y \approx yx^2.$$

Open problem

Classify all modular elements of the lattice **MON**.

An element a of a lattice L is called *modular* if

$$(\forall x, y \in L) \quad x \leq y \longrightarrow (a \vee x) \wedge y = (a \wedge y) \vee x.$$

Necessary condition of the modularity

Let V be a proper monoid variety that is a modular element of the lattice **MON**. Then V satisfies the identities

$$x^2 \approx x^3 \text{ and } x^2y \approx yx^2.$$

Open problem

Classify all modular elements of the lattice **MON**.

An element a of a lattice L is called *modular* if

$$(\forall x, y \in L) \quad x \leq y \longrightarrow (a \vee x) \wedge y = (a \wedge y) \vee x.$$

Necessary condition of the modularity

Let V be a proper monoid variety that is a modular element of the lattice **MON**. Then V satisfies the identities

$$x^2 \approx x^3 \text{ and } x^2y \approx yx^2.$$

Open problem

Classify all modular elements of the lattice **MON**.

Thank you for your attention!

Thank you for your attention!